



**Animal and Plant Health
Inspection Service**

4700 River Road
Riverdale, MD 20737
Voice 301-851-4100
Web: <http://www.aphis.usda.gov>

News Release

Contact:
Joelle Hayden (301) 851-4040
joelle.r.hayden@aphis.usda.gov

Ed Curlett (301) 851-4052
ed.c.curlett@aphis.usda.gov

Highly Pathogenic H5N8 Avian Influenza Confirmed in Commercial Turkey Flock in California

No immediate public health concern; detected strain not known to harm humans

WASHINGTON, Jan. 24, 2015— The United States Department of Agriculture's (USDA) Animal and Plant Health Inspection Service (APHIS) has confirmed the presence of highly pathogenic (HPAI) H5N8 avian influenza in a commercial turkey flock in Stanislaus County, California. This is the first finding of HPAI in commercial poultry during the ongoing disease incident in the Pacific Flyway. **No human cases of these avian influenza viruses have been detected in the United States, Canada, or internationally, and there continues to be no immediate public health concern.**

Samples from the flock, which experienced increased mortality, were tested at the California Animal Health & Food Safety Laboratory System (CAHFS) and the APHIS National Veterinary Services Laboratories in Ames, Iowa confirmed the finding. APHIS is partnering closely with the California Department of Food and Agriculture (CDFA), which has quarantined the facility. APHIS and CDFA have initiated an incident command response, and APHIS will assist CDFA in depopulating the remaining birds on the property to prevent the spread of the disease.

While birds from the affected flock will not enter the food chain, eating properly handled and cooked poultry and eggs is safe. Cooking poultry to an internal temperature of 165 °F kills bacteria and viruses.

Federal and State partners are working jointly on additional surveillance and testing in the nearby area, following existing avian influenza response plans. These plans also will include preventing the movement of risky animals or products out of the immediate area to prevent further disease spread. The United States has the strongest AI surveillance program in the world, and USDA is working with its partners to actively look for the disease in commercial poultry operations, live bird markets, and in migratory wild bird populations.

USDA will be notifying the World Organization for Animal Health (OIE) of this detection as part of USDA's ongoing reporting of all HPAI findings. USDA also continues to communicate with trading partners to encourage adherence to OIE standards and minimize trade impacts. OIE trade guidelines call on countries to base trade restrictions on sound science and, whenever possible, limit restrictions to those animals and animal products within a defined region that pose a risk of spreading disease of concern. .

These virus strains can travel in wild birds without them appearing sick. People should avoid contact with sick/dead poultry or wildlife. If contact occurs, wash your hands with soap and water and change clothing before having any contact with healthy domestic poultry and birds.

All bird owners, whether commercial producers or backyard enthusiasts, should continue to practice good biosecurity, prevent contact between their birds and wild birds, and to report sick birds or unusual bird deaths to State/Federal officials, either through your state veterinarian or through USDA's toll-free number at 1-866-536-7593. Additional information on biosecurity for backyard flocks can be found at healthybirds.aphis.usda.gov

Additional background:

Avian influenza (AI) is caused by an influenza type A virus which can infect poultry (such as chickens, turkeys, pheasants, quail, domestic ducks, geese, and guinea fowl) and is carried by free flying waterfowl such as ducks, geese and shorebirds. AI viruses are classified by a combination of two groups of proteins: hemagglutinin or "H" proteins, of which there are 16 (H1–H16), and neuraminidase or "N" proteins, of which there are 9 (N1–N9). Many different combinations of "H" and "N" proteins are possible. Each combination is considered a different subtype, and can be further broken down into different strains. AI viruses are further classified by their pathogenicity (low or high)—the ability of a particular virus strain to produce disease in domestic chickens.

The H5N8 virus originated in Asia and spread rapidly along wild bird migratory pathways during 2014, including the Pacific flyway. In the Pacific flyway, the H5N8 virus has mixed with North American avian influenza viruses, creating new mixed-origin viruses. These mixed-origin viruses contain the Asian-origin H5 part of the virus, which is highly pathogenic to poultry. The N parts of these viruses came from North American low pathogenic avian influenza viruses.

USDA has identified two mixed-origin viruses in the Pacific Flyway: the H5N2 virus and new H5N1 virus. The new H5N1 virus is **not** the same virus as the H5N1 virus found in Asia, Europe and Africa that has caused some human illness. Detailed analysis of the virus is underway in cooperation with the U.S. Centers for Disease Control and Prevention. None of these viruses have been identified in humans, nor are expected to pose a public health risk.

For more information about the ongoing avian influenza disease incident in the Pacific Flyway visit the [APHIS website](#). More information about avian influenza can be found on the [USDA avian influenza](#) page.

#